

Bactur[®]

CATERPILLAR
KILLER



4547

**Controls caterpillars on vegetables,
fruit and ornamentals.**

- Safe to handle
- Bee and plant safe - **BioGro** certified
- Nil withholding period

Grosafe[®] Bactur[®] Caterpillar Killer contains naturally occurring bacteria (*Bacillus thuringiensis*)

FOR BEST RESULTS:

- Apply when caterpillars are young, before plant damage occurs.
- Shake frequently during application.
- Spray both upper and lower surfaces of leaves if possible.
- Do not use when rain is forecast within 8 hours.
- Broaden insect spray control and enhance efficacy by adding **Grosafe[®] EnSpray 99[®] Spraying Oil** to mixture.



How to Use BACTUR® Caterpillar Killer

COMPATIBILITY: To achieve a broad spectrum pest kill add **Grosafe® EnSpray 99®** Spraying Oil at label rates, but not with products that are either alkaline in reaction or based on heavy metals (e.g. copper based products).

Mix **Grosafe® Bactur®** Caterpillar Killer mixture first with water then add the **Grosafe® EnSpray 99®** Spraying Oil at label rates.



Controls:



White butterfly caterpillar



Leaf roller caterpillar



Diamond backed moth caterpillar



Looper caterpillar



Tomato fruitworm

NOTE: Do not use in conjunction with foliar nutrients, liquid fertilisers or Bordeaux mixture. If mixing with cupric hydroxide, use immediately, do not allow to stand.

Wash fruit and vegetables well before eating.

DIRECTIONS: 1. Dilute sachet contents into 5 Litres of water. 2. Spray to the point of runoff. 3. Spray as recommended (see below).

Apples, Avocados, Berry Fruits, Citrus, Grapes, Kiwifruit, Pipfruit	Leaf roller caterpillars	Apply at 10 - 14 day intervals at first sign of caterpillars, applying at early, mid and late flowering. Ensure thorough coverage. Broaden control by using Grosafe® EnSpray 99® Spraying Oil at label rates.
Tomatoes	Tomato fruitworm, Looper caterpillar	Apply according to monitoring, or at first sign of attack. Repeat at 10 - 14 day intervals until harvest.
Vegetables Brassicas (cabbage brussel sprouts, broccoli, cauliflower)	White butterfly caterpillar, Diamond backed moth caterpillar	Apply at 10 - 14 day intervals at first sign of caterpillars, applying at early, mid and late flowering. Ensure thorough coverage.
Ornamentals, other fruit, other vegetables	White butterfly Caterpillar, Diamond back moth caterpillar	Apply every 7 - 10 days, as required until control is achieved.
Withholding period: Nil		



Grosafe garden

20 Jean Batten Drive, Mt Maunganui, 3116
PO Box 14 450 Tauranga 3143.
Ph: +64 7 572 2662 | info@grosafe.co.nz
www.grosafegarden.co.nz